

Measurement of longitudinal single-spin asymmetries for W^\pm boson production in polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV at STAR

Devika Gunarathne^{*†}

Temple University, Philadelphia, PA, USA

E-mail: devika@temple.edu

W^\pm boson production in longitudinally polarized $p + p$ collisions provides unique and clean access to the individual helicity polarizations of u/d quarks and anti-quarks. Due to maximal violation of parity, W bosons couple to left-handed quarks and right-handed anti-quarks and hence offer direct probes of their respective helicity distributions in the nucleon. These can be extracted from measured parity-violating longitudinal single-spin asymmetries, A_L , for $W^{+(-)}$ boson production as a function of the decay lepton (positron) pseudo-rapidity η . The STAR experiment is well equipped to measure A_L for W^\pm boson production for $|\eta| < 1$. The published STAR A_L results (2011 and 2012 data combined) have been used by several theoretical analyses suggesting a significant impact in constraining the helicity distributions of anti- u and anti- d quarks. In 2013 the STAR experiment has collected a large data sample of $\sim 250 \text{ pb}^{-1}$ which is more than 3 times larger than the total integrated luminosity in 2012, at $\sqrt{s} = 510$ GeV with an average beam polarization of $\sim 54\%$, comparable to run 2012. The status of the 2013 A_L analysis will be discussed along with an overview of future plans.

The XXIII International Workshop on Deep Inelastic Scattering and Related Subjects

April 27 - May 1, 2015

Southern Methodist University

Dallas, Texas 75275

*Speaker.

†for the STAR collaboration

1. Introduction

There has been steady progress over the past couple of decades in terms of understanding the spin structure of the nucleon, one of the fundamental questions in nuclear physics. In the 1980s, the spin of the proton was naively explained by the alignment of spins of the valence quarks. However, in our current understanding, the valence quarks, sea quarks, gluons and their possible orbital angular momentum are all expected to contribute to the overall spin of the proton. Despite this significant progress, our understanding of the individual polarizations of quarks and antiquarks is not yet complete.

According to the spin sum rule of a longitudinally polarized proton [2], the spin of the proton can be written in terms of its contributions from the intrinsic quark and antiquark polarization, intrinsic gluon polarization and their possible orbital angular momentum. Polarized inclusive deep-inelastic scattering (DIS) experiments were able to strongly constrain the total quark contribution to the proton spin [1]. However, DIS experiments were not sensitive to the flavor separated individual quark spin contributions. These were then measured by polarized semi inclusive DIS experiments (SIDIS), but relatively large uncertainties were observed in the extracted helicity-dependent parton distribution functions (PDF) [1] due to the use of hadronic fragmentation functions in reconstructing the kinematics of the final state particles.

The production of W^\pm bosons in longitudinally polarized $p + p$ collisions at RHIC provide an unique and powerful tool to probe the individual helicity PDFs of light quarks and anti-quarks in the proton. Due to the maximal violation of parity during the production, $W^{-(+)}$ bosons couple to the left-handed quarks and right-handed anti-quarks and hence offer direct probes of their respective helicity distributions in the nucleon. These distributions can be extracted by measuring the parity-violating longitudinal single-spin asymmetry, A_L , as a function of the decay electron (positron) pseudo-rapidity, η_e . The longitudinal single-spin asymmetry is defined as $A_L = (\sigma_+ - \sigma_-) / (\sigma_+ + \sigma_-)$, where $\sigma_{+(-)}$ is the cross section when the helicity of the polarized proton beam is positive (negative). At leading order, $W^+ A_L$ is directly related to polarized anti d and u quark distributions ($\Delta\bar{d}$, Δu) while $W^- A_L$ is directly related to polarized anti u and d quark distributions ($\Delta\bar{u}$, Δd) [3].

The results of the single-spin asymmetries for W^\pm boson production in longitudinally polarized $p + p$ collisions from the 2011 and 2012 STAR running periods are presented. The integrated luminosity of the data set collected during these two years were 9 and 77 pb^{-1} , with an average beam polarization of 49% and 56%, respectively.

2. Analysis

The STAR experiment is well equipped to measure A_L for W^\pm boson production within a pseudorapidity range of $|\eta| < 1$. W^\pm bosons are detected via their $W^\pm \rightarrow e^\pm \nu$ decay channels. A subsystem of the STAR detector, the Time Projection Chamber (TPC) is used to measure the transverse momentum (p_T) of decay electrons and positrons and to separate their charge sign. Two other subsystems, Barrel and Endcap Electromagnetic Calorimeters (BEMC, EEMC) are used to measure the energy of decay leptons. A well developed algorithm is used to identify and reconstruct W^\pm candidate events by reducing large QCD type background events. In this algorithm, various

cuts are designed at each level of the selection process based on the kinematics and topological differences between the electroweak process of interest and QCD processes. For example, tracks associated with W^\pm candidate events can be identified as isolated tracks in the TPC that point to an isolated EMC cluster in the calorimeter, where as for QCD type events have several TPC tracks point to several EMC clusters. In contrast to QCD background events, large opposite missing transverse energy can be observed in $W^\pm \rightarrow e^\pm \nu$ decay, due to undetected neutrinos. This leads to a large imbalance in the vector p_T sum of all reconstructed final-state objects in W candidate events, which is expressed as $\vec{p}_T^{balance}$ in equation 2.1. Here \vec{p}_T^{jets} is determined using the anti- k_T algorithm [3]. A cone of radius of 0.7 in $\eta - \phi$ space is centered around the candidate lepton, and the p_T from all reconstructed jets outside of the cone are included. The \vec{p}_T^e is the p_T of the candidate lepton. A strong correlation can be observed between E_T and scaler quantity, signed p_T balance, which is defined in equation 2.2. This can be seen clearly in Figure 1 c) which shows the MC results simulating $W^\pm \rightarrow e\nu$ decays. After initial requirement of reconstructed TPC tracks to have $p_T > 10 GeV$, tracks are matched to a 2×2 EMC cluster with transverse energy $E_T > 12 GeV$. W candidate events are then isolated using the large opposite missing energy requirement. This is done by requiring that the E_T fraction of the $2 \times 2 / 4 \times 4$ tower clusters is larger than 95%. The final requirement, which is based on the large imbalance in the vector p_T sum mentioned above, is signed- p_T balance to be greater than 14 GeV, which indicated by the red line in Figure 1 a). After all the selection cuts have been applied, the characteristic Jacobian peak in the E_T distribution for mid-rapidity W^\pm candidate events can be observed near half of the W^\pm mass, as shown in Figure 1 (a).

$$\vec{p}_T^{balance} = \vec{p}_T^e + \sum_{\Delta R > 0.7} \vec{p}_T^{jets} \quad (2.1)$$

$$signed\ p_T\ balance = \frac{(\vec{p}_T^e) \cdot \vec{p}_T^{balance}}{|\vec{p}_T^e|} \quad (2.2)$$




Figure 1: Candidate E_T^e distribution from the data after various selection cuts (a), Signed p_T -balance vs E_T^e for data (b) and $W \rightarrow e\nu$ MC (c)

The Charge separated W^\pm yields from the 2011 and 2012 data sets as a function of E_T^e are shown in Figure 2 for different η bins, along with the estimated residual background contributions from $W^\pm \rightarrow \tau^\pm \nu_\tau$, $Z/\gamma^* \rightarrow e^+e^-$ electroweak processes and QCD processes. Relatively small electroweak background contributions are estimated from Monte-Carlo (MC) simulation, with PYTHIA 4.622 generated events passing through the STAR GEANT model and embedded in to STAR zero-bias triggered events. Despite a significant reduction of QCD background events during the selection process, a certain amount is still present in the signal region. This contribution is primarily originated from events which satisfy candidate W^\pm isolation cuts but contain jets which escape the detection outside the STAR acceptance. Two procedures referred as "Second EEMC" and "Data-driven QCD" [5], are used to estimated the background associate with the acceptance ranges $-2 < \eta < -1.09$ and $|\eta| > 2$. At forward rapidity, ($1 < \eta_e < 1.4$), the W selection criteria used is similar to that of mid rapidity. The Background estimation is improved using additional Endcap Shower Maximum Detector (ESMD). More details are described in [4].




Figure 2: E_T^e distribution of W^- (top) and W^+ (bottom) candidate events (black), various background contributions and sum of backgrounds and $W \rightarrow e \nu$ MC signal (red-dashed).

3. Results

To properly account for the low statistics in the 2011 data set, a profile likelihood method was used to extract the spin asymmetry results from the combined 2011 and 2012 data sets. Two likelihood functions L_{year1} and L_{year2} are defined for each of the 2011 and 2012 data sets respectively. The Asymmetry results, central values and confidence intervals are extracted from the product of

the likelihood function, $L_{2011} \times L_{2012}$. More details are described in [4]. The W^\pm single-spin asymmetry results measured for e^\pm with $25 < E_T^e < 50 \text{ GeV}$ are shown in Figure 3 as the function of decayed e^\pm pseudorapidity, η_e in comparison to theoretical predictions based on DSSV08 [6] and LSS10 [7] helicity-dependent PDF sets, using both CHE (next-to-leading order) [3] and RHICBOS (fully resummed) frameworks [8]. The measured $A_L^{W^-}$ is larger than the central value of the theoretical predictions. The enhancement at large negative η_e , in particular is sensitive to polarized anti u quark distribution, $\Delta\bar{u}$. $A_L^{W^+}$ is negative as expected and consistent with theoretical predictions. The systematic uncertainties for $A_L^{W^\pm}$ are well under control for pseudorapidity range $|\eta_e| < 1.4$. STAR 2012 preliminary $A_L^{W^\pm}$ results [1] are included in the DSSV++ global analysis [9] from the DSSV group and recent NNPDF [10] global analysis. Both analyses shows that the STAR $W A_L$ results provide a significant constraint on anti u ($\Delta\bar{u}$) and anti d ($\Delta\bar{d}$) quark polarizations.




Figure 3: Longitudinal single-spin asymmetries for W^\pm production as a function of lepton pseudorapidity, η_e in comparison to theory predictions.

4. Outlook

In 2013, the STAR experiment collected a large data sample of $\sim 250 \text{ pb}^{-1}$, which is more than 3 times larger than the total integrated luminosity in 2012, at $\sqrt{s} = 510 \text{ GeV}$ with an average beam polarization of $\sim 54\%$. The high luminosity data collected in 2013 requires proper calibration of

all the subsystems used in the analysis. As the charge sign reconstruction of e^+ and e^- is based on the bending of TPC tracks in the presence of an axial magnetic field, the calibration of the TPC is crucial in W analysis. With the large pile up accumulated in the TPC due to high luminosity $p + p$ collisions in 2013, this is extremely challenging. Despite this challenging environment the completed TPC calibration for STAR 2013 data set indicate a clear charge separation between e^+ and e^- . The calibration of the other crucial subsystem for mid-rapidity W analysis, the BEMC, is currently in progress. Forward Gem Tracker (FGT) was fully installed at STAR in year 2013, which covers the acceptance between $1 < \eta < 2$ in the forward region. The FGT will be used as the tracking device for W analysis in the forward pseudorapidity region at STAR. This enhances the sensitivity to \bar{u} and \bar{d} quark polarizations. Figure 4 shows the projected uncertainties for the W^\pm asymmetries estimated from the 2013 data set.




Figure 4: Projected uncertainties for the longitudinal signal-spin asymmetries as a function of lepton pseudorapidity, η_e for W^\pm production from the 2013 data set.

Higher precision results are expected from the STAR 2013 $W A_L$ analysis to improve constraints on the sea quarks helicity-dependent PDFs.

References

- [1] D. de Florian, R. Sassor, M. Stratmann, and W. Vogelsang, Extraction of Spin-Dependent Parton Densities and Their Uncertainties, *Phys. Rev.* **D80**, 034030 (2009).
- [2] R. Jaffe and A. Manohar, The G(1) Problem: Fact and Fantasy on the Spin of the Proton, *Nucl. Phys.* **B337**, 509 (1990), revised version.
- [3] D. de Florian and W. Vogelsang, Helicity parton distributions from spin asymmetries in W-boson production at RHIC, *Phys. Rev.* **D81**, 094020 (2010).
- [4] STAR Collaboration, L. Adamczyk *et al.*, *Phys. Rev. Lett.* **113**, 072301 (2014).
- [5] STAR Collaboration, L. Adamczyk *et al.*, *Phys. Rev.* **D85**, 092010 (2012).
- [6] D. de Florian, R. Sassor, M. Stratmann, and W. Vogelsang, *Phys. Rev. Lett.* **101**, 072001 (2008).
- [7] E. Leader, A. V. Sidorov, and D. B. Stamenov, Determination of polarized parton densities from a QCD analysis of inclusive and semi-inclusive deep inelastic scattering data, *Phys. Rev.* **D82**, 114018 (2010).
- [8] P. M. Nadolsky and C. Yuan, *Nucl. Phys.* **B666**, 31 (2003)
- [9] E. Aschenauer *et al.*, (2013), arXiv:1304.0079 [nucl-ex].
- [10] E.R. Nocera, arXiv:1403:0440 [hep-ph] (2014).